# CONCEPTUAL AND NUMERICAL MODEL OF INFILTRATION FOR THE YUCCA MOUNTAIN AREA, NEVADA

U.S. GEOLOGICAL SURVEY

Water Resources Investigation Report ##-###

DRAFT

Prepared in cooperation with the NEVADA OPERATIONS OFFICE U.S. DEPARTMENT OF ENERGY, under Interagency Agreement DE-AI08-92NV10874

Dept. Seal **Version** September 20, 1996

INFIL\_R1.WPD

# CONCEPTUAL AND NUMERICAL MODEL OF INFILTRATION FOR THE YUCCA MOUNTAIN AREA, NEVADA

by Alan L. Flint, Joseph A. Hevesi and Lorraine E. Flint

U.S. GEOLOGICAL SURVEY

Water Resources Investigation Report ##-###

Prepared in cooperation with the NEVADA OPERATIONS OFFICE U.S. DEPARTMENT OF ENERGY, under Interagency Agreement DE-AI08-92NV10874

Denver, Colorado 1996

#### U.S. DEPARTMENT OF THE INTERIOR

### **BRUCE BABBITT, Secretary**

U.S. GEOLOGICAL SURVEY

Gordon P. Eaton, Director

The use of trade, product, industry, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government

For additional information write to:

Chief, Earth Sciences Investigations Program Yucca Mountain Project Branch U.S. Geological Survey Box 25046, MS 421 Denver Federal Center Denver, CO 80225 Copies of this report can be purchased from:

U.S. Geological Survey
Earth Science Information Center
Open-File Reports Section
Box 25286, MS 517
Denver Federal Center
Denver, CO 80225

## **CONTENTS**

ABSTRACT1
INTRODUCTION
Background
Purpose and Objectives
Previous Work
Previous Work
PROCESSES, THE PHYSICS OF INFILTRATION
Field Water Balance
Precipitation
Run-on / Runoff
Intiltration \
Evapotranspiration
Redistribution
Flow at the soil-bedrock interface
Net infiltration
25
STUDY AREA DESCRIPTION
Geographic and Physicographic Setting
Geographic and Physiographic Setting
Climate
Present
Present
Future
Geology
Distribution
Properties
Watershed Characteristics
Defined watersheds
Topographic surfaces
30
Distribution
Properties
Vegetation
•
DATA COLLECTION / DATA ANALYSIS
Frecipitation
ASSIST OF A REGISTRATE AND A SECOND OF THE S
minutation
Evaportaispiration
TCUISTIUMQII
Neutron hole moisture profile analysis
,

INFIL\_R1.WPD

iy

September 27, 1996

Geochemistry analysis	<i>-</i> 2
Flow at the soil-bedrock interface  Net infiltration	נט
Net infiltration	66
Net infiltration	68
NUMERICAL MODEL	60
4 Coophanon	
Texts-on/Addition	
Infiltration  Evanotranspiration	/2
Evapotranspiration	74
Evapotranspiration Solar radiation	74
Solar radiation	74
arabadisea x resitey-1 avior equinion	
1464 HAMINGHOTE A	
Stochastic Simulations of Infiltration	15
	5
ASSUMPTIONS AND LIMITATIONS	
ASSUMPTIONS AND LIMITATIONS	8
FUTURE WORK	9
SUMMARY 90	^
	U
Appendix I All Yucca Mountain Project data referred to in the	
Appendix I All Yucca Mountain Project data referred to in this report, listed by data tracking	
number	4
Appendix II Neutron borehole locations and information used in numerical model. $\dots$ 107	7
1000 model	′
Appendix III Program MARKOV 108	
	3
Appendix IV Program PRISIM	
Appendix IV Program PPTSIM	•
Appendix V Program INFIL	7
	I

INFIL\_R1.WPD

September 27, 1996

### **FIGURES**

1. Location of a) North American Deserts a	and the original 3 s	sites for characterization, and b)
physiographic provinces		
2. Yucca Mountain regional study area with	h Death Valley Gr	ound-Water Unit and infiltration
study boundary		
3. Field scale schematic of infiltration		
4. Diagrammatic water balance components	s using notation de	escribed in equation 1
5. Conceptual model of the energy balance	equation for evapo	otranspiration.
6. a)Water content and water potential at the	e soil-bedrock inte	rface at a denth of 2.1 m in
borehole USW UZ-N52. The horizo		
um open fracture and a filled fracture	e, and b) associated	flux calculated as the hydraulic
conductivity of the fracture times a u	mit gradient	25
7. Infiltration study area and modeling doma	ain with original si	ite-scale model houndary (Wittwer
and others, 1992) and potential repos	sitory location	27
8. Historical climate change estimates from	a) SPECMAP stuc	lies with a hypothetical transfer to
precipitation, b) high-frequency fluct	uations in climate	change from the GRIP studies and
c) recharge estimated from climate ch	nange cycles using	the Maxey-Eakin method 30
9. Predominate direction of weather patterns	s and seasonal sour	ces of moisture across the
southwestern U.S.	******	21
10. Map of area with general geologic design	nations (after Saw	ver and others 1994)
11. Saturated hydraulic conductivity of the st	urficial bedrock ur	nits
12. Map of major watersheds at Yucca Mour	ntain	27
13. Map of soil thickness designated in 4 cate	egories, $0 - < 0.5$	0.5 = < 3  m  3 = < 6  m  and  5  m
		$0.5 \times 5 \text{ m}, 5 \times 6 \text{ m}, \text{ and } \geq 6 \text{ m}.$
14. Map of surficial soils		41
15. Relationship of a) soil texture to saturated	d hydraulic conduc	tivity using Comphell (1995) s
soil without rock fragments and modif	fied to correct for a	percent rock from sets and 11)
modeled versus measured conductivity	V	recommendation magnifiles, and d)
	, <u></u>	
INFIL_R1.WPD	vi	September 20, 1007

16.	Moisture retention curves calculated from measured soil texture (Campbell, 1985) compared
	to measured data for soil model units a) 1, b) 2, and c) 4, and texture models for d) all soi
	model units
17.	Map of vegetation associations
18.	Rate of rainfall for a) a summer thunderstorm and b) winter precipitation47
19.	Estimated daily precipitation for Yucca Mountain using 15 years of measured data simulated
	at 90 neutron boreholes
20.	Probability of occurrence of daily precipitation at Yucca Mountain using 3rd order Markov
	chains. A 4-day window was used and 1 equals precipitation
21.	Data and fitted model for the cumulative probability distribution of precipitation quantity at
	station 4JA a) for the month of February and b) for the month of July
22.	Probability distribution of precipitation quantity for a) precipitation data from Area 12 and
	b) precipitation data from Yucca Mountain
23.	Hydrograph for Pagany Wash and Wren Wash, March 11, 1995
24.	Map of maximum infiltration capacity into initially air-dry soil
25.	Map of minimum infiltration capacity
26.	Illustration of the change in infiltration capacity for soil model unit 5 with time, related to
	precipitation rate and minimum infiltration capacity58
27.	Map of field capacity of surficial soils
28.	Field storage capacity of soils calculated as field capacity multiplied by soil depth 59
	Measured evaporation pan data and calculated potential evapotranspiration using energy
	balance for 1991-1993
30.	Average depth of penetration of the wetting front for years 1990-1993 and soil depth for 34
	boreholes (Flint and Flint, 1995)
31.	Changes in water content over time and depth for borehole USW UZ-N1 62
32.	Changes in water content over time and depth for borehole USW UZ-N15 63
33.	Saturation, porosity and <sup>36</sup> Cl/Cl ratios for boreholes USW UZ-N15, USW UZ-N16, USW
	UZ-N17 and USW UZ-N11. 36Cl data is indicated for depth ranges as between pairs of
	points
`	D.I. W.D.D.

INFI	_R1.WPD viii September 20, 1996	
48.	Alternative conceptual model of infiltration numerically modeled assuming all infiltration	is
	Close-up view of the potential repository area in figure 46	
	resulting in an average of 4.5 mm/year flux over the modeling domain	
46.	Conceptual model of infiltration numerically modeled using average annual precipitation	
	precipitation values of a) 165 mm/year and b) 205 mm/year	86
45.	nfiltration simulated using stochastic precipitation simulations with average annual	
	Eakin method and calculated from neutron borehole data	85
44.	Precipitation versus net infiltration, modeled 100-year simulation, estimated using Maxey-	-
	all boreholes. Linear regressions predicting values for each data set are shown	84
	factor and with 30 percent channel enhancement factor, and mean yearly infiltration for	г
43.	Precipitation versus infiltration modeled for each borehole with no channel enhancement	
	data	84
	enhancement factor compared to simulated average annual precipitation for 15 years of	
42.	recipitation modeled with no channel enhancement factor and with 30 percent channel	
	USW UZ-N63	84
41.	Calibration of modeled net infiltration using neutron boreholes a) USW UZ-N50 and b)	
	simulations of precipitation from a) Yucca Mountain and b) Area 12	83
40.	Simulation of mean net infiltration for neutron boreholes using 100-year stochastic	
	years using the YM stochastic rainfall model	81
39.	Volumetric water content simulated at one location in Split Wash for a) 15 years and b) 1	00
	precipitation simulated at 90 neutron boreholes	76
38.	Simulated daily evapotranspiration for Yucca Mountain using 15 years of measured	
37.	Average annual precipitation distributed spatially using an elevation correlation	
	12	71
36.	One hundred-year stochastic simulations of precipitation at a) Yucca Mountain and b) Are	a
	measured at 4 depths for 1995 and b) used to calculate flux.	68
35.	Water potential measurements near borehole USW UZ-N15 using heat dissipation probes,	, a)
	UZ-N55. <sup>36</sup> Cl data is indicated over a depth range	66
34.	Saturation, porosity and 36Cl/Cl ratios for boreholes USW UZ-N53, USW UZ-N54 and US	SW

	through the channels, and assuming an average of 10 mm/yr, resulting in an average flux
	of 0.5 mm/yr over the modeling domain
TA	ABLES
1.	Generalized lithostratigraphy of Yucca Mountain area and ages
2.	Properties of rock matrix and fractures for all lithostratigraphic units exposed in the study
	area.
	36
3.	Surficial soil units recombined from mapped surficial units
4.	Summary of soil properties
5.	Plant species most commonly found at Yucca Mountain
6.	Model parameters and cumulative probability for various minimum amounts of daily
	precipitation for stations 4JA and Area 12 calculated for months
7.	Infiltration fluxes estimated by the chloride mass balance method (adapted from Fabryka-
	Martin and others, 1994) with a comparison of point and areally averaged estimates of net
	infiltration

#### **CONVERSION FACTORS**

Multiply	Ву	To Obtain
millimeter (mm)	0.03937	inch (in)
meter (m)	3.28	foot (ft)
watt (w)	0.2388	calorie per second (C/s)
kilopascal (KPa)	0.01	bar
kilometer (km)	0.6214	mile (mi)
square kilometer (km²)	0.386	square mile (mi <sup>2</sup> )
millimeter per year (mm/yr)	0.00328	foot per year (ft/yr)
cubic centimeter (cm³)	3.531 x 10 <sup>-5</sup>	cubic foot (ft³)
cubic centimeter per second (cm³/s)	3.531 x 10 <sup>-5</sup>	cubic foot per second (ft³/s)

Degree Celsius (°C) may be converted to degree Fahrenheit (°F) by using the following equation:

$$^{\circ}F = 9/5 \, ^{\circ}C + 32.$$